The extracellular electron transfer capabilities of rumen microbes are stimulated by conductive nanomaterials

<u>Abimael Ortiz-Chura</u>, Jeremy Tournayre, Aurelie Dehornoy, Milka Popova and Diego Morgavi

Université Clermont Auvergne, INRAE, VetAgro Sup, UMR 1213 Herbivores Unit, Saint-Gènes-Champanelle, France






> Strategies to explore Extracellular Electron Transfer (EET)

Rumen microbes capable to EET via DIET/redox protein remain undetected and underexplored

- ➤ Conductive materials (CMs) increase EET
 - ➤ Can promote biological processes
 - ➤ ↑ methane production in some ecosystems
- ➤ Carbon-based materials
 - ➤ Activated charcoal and graphene nanoplatelets
 - ➤ Provide an electron-conducting surface
- ➤ Mineral-based materials
 - ightharpoonup Magnetite [$Fe^{2+}Fe_2^{3+}O_4$; Iron(II,III) oxide]
 - > Acts as an electron conduit





Hypothesis

- ➤ Incubation of CMs, graphene and magnetite, in the rumen would enrich microbial communities associated with EET that induce shifts in fermentation parameters
- Does it change fermentation parameters? Methane?
- Which microbes could be associated with this presumed EET mechanism in the rumen?

Objective

➤ To test the effect of CMs on rumen fermentation and on microbial communities



METHODS

In vitro experiments

Rumen inoculum: Sheep

Treatments:

Control: Substrate alone

Graphene (GPH) 40 mg/bottle

Magnetite (MGN) 40 mg/bottle

Substrates: Solid or soluble

In sacco experiment

Rumen incubation: Sheep


Treatments:


Control: polymer PDMS alone


PDMS + Graphene (GPH)

PDMS + Magnetite (MGN)

Incubation time: 1d, 7d and 28d







Isolation



p. 4

> RESULTS: Fermentation


In vitro batch culture (48h)

Items	CTR	GPH 5%	GPH 10%	MGN 5%	MGN 10%	SEM	P-value
Gas production (mL)	156,0	154,0	157,0	158,0	155,0	2.24	0.542
Methane production (mL)	19.3	19.5	20.3	20.3	20.1	0.43	0.131
рН	6.84	6.81	6.81	6.78	6.77	0.03	0.321
Total VFA	107.1	100.3	96.7	98.7	96.6	3.42	0.053
Acetate	65.3	65.4	65,0	64.3	64,0	0.76	0.319
Propionate	18,0	17.7	18.6	18.5	18.2	0.48	0.342
Butyrate	7.7	7.8	7.3	7.7	8.1	0.35	0.271
Isobutyrate	2.1	2.1	2.1	2.1	2.1	0.02	0.268
Valerate	3.6	3.6	3.6	3.6	3.6	0.07	0.764
Isovalerate	3.1	3.2	3.1	3.4	3.7	0.21	0.072
Caproate	0.2	0.3	0.2	0.3	0.4	0.04	0.060
A/P	3.7	3.7	3.5	3.5	3.5	0.14	0.392

No major differences between treatments for any fermentative metabolites

CMs tend to increase methane production (~5%)






INRAO

In sacco experiment

> RESULTS: Fermentation


Consecutive batch culture



CMs increased methane production

→ Total VFA and acetate

□ propionate






RESULTS: Microbiota Consecutive batch culture - Soluble substrate

inoculum and CTR

WUniFrac

S3 to S10

2025-06-02 RMG



p. 8

Summary

CMs Increased methane production
How? Enriched pathways/genes?

➤ Shift in microbes associated to conductive materials

suggests the presence of EET in the rumen

Correlation is not causation!

- EET and mechanisms involved should be validated with additional tools
- Its extent in the rumen should be assessed as it could be another mechanism to consider for modulating methanogenesis




Limitation and future directions


Next steps:

- Functional analysis (genes/pathways)
- Isolation
- ✓ A *Methanomicrobium mobile* strain was isolated
- Co-culture

Geobacter a well know DIET bacteria

G. Metallireducens <=> M. mobile?





Acknowledgments



We acknowledge the support by the EU Horizon 2020 research and innovation program under grant agreements101000213-HoloRuminant.



Thank you for your attention!

